

Rodney
Undetectable
since 2008

I didn't come this far without staying informed.
Today, I want to find out more about
**staying undetectable* on a complete HIV treatment
with fewer medicines in 1 small pill.**



What is JULUCA?

JULUCA is a prescription medicine that is used without other antiretroviral medicines to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults to replace their current anti-HIV-1 medicines when their healthcare provider determines that they meet certain requirements. HIV-1 is the virus that causes Acquired Immune Deficiency Syndrome (AIDS). It is not known if JULUCA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take JULUCA if you:

- have ever had an allergic reaction to a medicine that contains dolutegravir or rilpivirine.

*Undetectable means keeping the amount of HIV-1 in the blood at very low levels (less than 50 copies per mL). Your results may vary.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.

Please see Important Facts about JULUCA on page 27.

Every person featured in this brochure is a real patient diagnosed with HIV-1. They have all been compensated for their time by ViiV Healthcare.

I'm not the same person I was
when I was first diagnosed with HIV.
Should I be taking the same HIV treatment?

Tiera
Undetectable since 2010

IMPORTANT SAFETY INFORMATION (cont'd)

Do not take JULUCA if you: (cont'd)

- are taking any of the following medicines:
 - dofetilide
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifampin
 - rifapentine
 - proton pump inhibitors (including esomeprazole, lansoprazole, omeprazole, pantoprazole sodium, rabeprazole)
 - St. John's wort (*Hypericum perforatum*)
 - more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.

Hello, Undetectable you.

JULUCA is a next step in HIV treatment for adults who have been undetectable* for at least six months.

JULUCA is a once-daily, complete HIV-1 regimen that combines 2 medicines in just 1 small pill.

2 medicines. 1 pill.

dolutegravir  rilpivirine 

2 things to know about JULUCA:

1.

It may keep you undetectable* with only 2 medicines in 1 pill.

2.

It's 1 small pill, taken once per day with a meal—in fact, it's the smallest† single-pill HIV-1 treatment available.

*Undetectable means keeping the amount of HIV-1 in the blood at very low levels (less than 50 copies per mL).

†Pill size is not intended to convey efficacy, safety, or treatment adherence. Approximate pill size is 14 mm x 7 mm.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA?

JULUCA can cause serious side effects, including:

- **Severe skin rash and allergic reactions.** Call your healthcare provider right away if you develop a rash with JULUCA. **Stop taking JULUCA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - blisters or sores in mouth
 - blisters or peeling of the skin
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue
 - problems breathing

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.

Now that you're undetectable,* have you thought about what's next?

There are many HIV treatment options available. Do you know how many medicines are in your HIV regimen?



➤ Ask your doctor if JULUCA could be a next step for you.

*Undetectable for at least six months. Undetectable means keeping the amount of HIV-1 in the blood at very low levels (less than 50 copies per mL).

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA? (cont'd)

JULUCA can cause serious side effects, including: (cont'd)

- **Liver problems.** People with a history of hepatitis B or C virus who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during treatment with JULUCA. Liver problems, including liver failure, have also happened in people without history of liver disease or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or “tea-colored” urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain, aching, or tenderness on the right side of your stomach area

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.

Alex
Undetectable
since 2011

**I went through a lot to get to undetectable,
and reaching it was one of the
greatest achievements of my life.**

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA? (cont'd)
JULUCA can cause serious side effects,
including: (cont'd)

- **Depression or mood changes. Tell your healthcare provider right away or get medical help if you have any of the following symptoms:**
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself
- **The most common side effects of JULUCA include:**
 - diarrhea
 - headache

These are not all the possible side effects of JULUCA. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider right away if you have any new symptoms while taking JULUCA.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see Important Facts about JULUCA on page 27.



 **Juluca**
dolutegravir 50 mg/
rilpivirine 25 mg tablets

**I've seen and done a lot over the years.
I've even learned I can stay undetectable
with fewer medicines in my HIV pill.**

Warren
Undetectable since 2010

IMPORTANT SAFETY INFORMATION (cont'd)

Before you take JULUCA, tell your healthcare provider if you:

- have ever had a severe skin rash or an allergic reaction to medicines that contain dolutegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had a mental health problem.
- are pregnant or plan to become pregnant. One of the medicines in JULUCA, called dolutegravir, may harm your unborn baby.
 - Your healthcare provider may prescribe a different medicine than JULUCA if you are planning to become pregnant or if pregnancy is confirmed during the first 12 weeks of pregnancy
 - If you can become pregnant, your healthcare provider will perform a pregnancy test before you start treatment with JULUCA, and you should consistently use effective birth control (contraception) during treatment with JULUCA
 - Tell your healthcare provider right away if you are planning to become pregnant, you become pregnant, or think you may be pregnant during treatment with JULUCA

**Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see Important Facts about JULUCA on page 27.**

Taking JULUCA.

Understanding the “how” and “when” of taking JULUCA is important. Be sure to keep these considerations in mind:



Not actual pill size.



Take JULUCA every day with a meal— exactly as your doctor tells you.

Taking JULUCA with a meal is important to help get the right amount of medicine in your body. A protein drink alone does not replace a meal.



Take JULUCA around the same time every day (or night).

Choose whatever time works best with your daily routine.



Take JULUCA as a replacement for your current regimen.

JULUCA is a complete HIV-1 regimen, so it isn't an addition to your current treatment, and shouldn't be taken with any other HIV-1 medicines.



Take JULUCA exactly as prescribed.

Do not miss a dose, change your dose, or stop taking JULUCA without talking to your doctor. Stay under a healthcare provider's care when taking JULUCA.

Some prescription and over-the-counter medicines interact with JULUCA and need to be taken at a different time from when you take JULUCA. See page 13 for more information.

Remember, it's important to keep your overall health in mind when considering HIV treatment options. Your doctor is a good source of information. Be sure to talk with him or her if you have questions.

➤ [Read our Doctor Discussion Guide on page 14.](#)

IMPORTANT SAFETY INFORMATION (cont'd)

Before you take JULUCA, tell your healthcare provider if you: (cont'd)

- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take JULUCA.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if JULUCA can pass to your baby in your breast milk.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.



 **Juluca**
dolutegravir 50 mg/
rilpivirine 25 mg tablets

The years have taught me how important
it is to learn more about my treatment options.
I'm ready to talk to my doctor about staying
undetectable with fewer medicines.

Ruben
Undetectable since 2009

IMPORTANT SAFETY INFORMATION (cont'd)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Some medicines interact with JULUCA. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with JULUCA.
- Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take JULUCA with other medicines.


How to take JULUCA?

- **Take JULUCA 1 time a day exactly as your healthcare provider tells you.**
- **Always take JULUCA with a meal.** A protein drink alone does not replace a meal.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.

Please see [Important Facts](#) about JULUCA on page 27.

Clinical trials.



This study included two groups of patients, those who took JULUCA for 100 weeks and those who switched from their current stable regimen‡ to JULUCA at week 52.

Patients who entered these studies were undetectable† and on their current stable regimen for at least six months. They could not participate if they had liver problems or active hepatitis B, were likely to need treatment for hepatitis C during the trials, had failed a prior HIV treatment, or had suspected resistance to the components of JULUCA (dolutegravir or rilpivirine).



*Both patients and their healthcare providers knew which medicines patients were taking.

†Undetectable means the amount of HIV-1 in the patients' blood was at very low levels (less than 50 copies per mL).

‡Regimens contained 2 nucleoside reverse transcriptase inhibitors (NRTIs) plus either an integrase strand transfer inhibitor (INSTI), a non-nucleoside reverse transcriptase inhibitor (NNRTI), or a protease inhibitor (PI).

IMPORTANT SAFETY INFORMATION (cont'd)

Do not take JULUCA if you:

- have ever had an allergic reaction to a medicine that contains dolutegravir or rilpivirine.
- are taking any of the following medicines:
 - dofetilide
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifampin
 - rifapentine
 - proton pump inhibitors (including esomeprazole, lansoprazole, omeprazole, pantoprazole sodium, rabeprazole)
 - St. John's wort (*Hypericum perforatum*)
 - more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.

The results are in.



In these studies, most people who were already undetectable* for at least six months stayed undetectable when they switched to JULUCA.†



AT 48 WEEKS

1 pill with 2 medicines was as effective at keeping people undetectable* as their previous 3- or 4-drug regimens.‡

95% of people who switched to JULUCA stayed undetectable*

95% of people who stayed on their current HIV regimen stayed undetectable*

AT 100 WEEKS

89% of people who took JULUCA for 100 weeks stayed undetectable*

93% of people who switched to JULUCA from week 52 to week 100 stayed undetectable*

*Undetectable means the amount of HIV-1 in the patients' blood was at very low levels (less than 50 copies per mL).

†Your results may vary.

‡Regimens contained 2 nucleoside reverse transcriptase inhibitors (NRTIs) plus either an integrase strand transfer inhibitor (INSTI), a non-nucleoside reverse transcriptase inhibitor (NNRTI), or a protease inhibitor (PI).

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA?

JULUCA can cause serious side effects, including:

- **Severe skin rash and allergic reactions.** Call your healthcare provider right away if you develop a rash with JULUCA. **Stop taking JULUCA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - blisters or sores in mouth
 - blisters or peeling of the skin
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue
 - problems breathing

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25. Please see [Important Facts](#) about JULUCA on page 27.

What can you expect?

You may experience some side effects when switching medications.

The most common side effects reported by people in these studies are*:

SWITCHED TO JULUCA AT THE START OF THE TRIALS

Common Side Effects	TAKING JULUCA	
	Weeks 1–48	Weeks 1–100
Diarrhea	2%	1%
Headache	2%	2%
Nausea	1%	2%

CONTINUED CURRENT HIV REGIMEN AT THE START OF THE TRIALS

Common Side Effects	TAKING CURRENT HIV REGIMEN	SWITCHED TO JULUCA
	Weeks 1–48	Weeks 52–100
Diarrhea	<1%	1%
Headache	0%	2%
Nausea	0%	1%

You should always talk to your doctor if you experience any side effect. Your results may vary.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

*Your results may vary.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA? (cont'd)

JULUCA can cause serious side effects, including: (cont'd)

- **Liver problems.** People with a history of hepatitis B or C virus who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during treatment with JULUCA. Liver problems, including liver failure, have also happened in people without history of liver disease or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or “tea-colored” urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain, aching, or tenderness on the right side of your stomach area

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25. Please see [Important Facts](#) about JULUCA on page 27.

What can you expect?

The discontinuation rates reported by people in these studies are*:

SWITCHED TO JULUCA AT THE START OF THE TRIALS

Discontinuation	TAKING JULUCA	
	Weeks 1–48	Weeks 1–100
People who left the study due to side effects	4%	7%

CONTINUED CURRENT HIV REGIMEN AT THE START OF THE TRIALS

Discontinuation	TAKING CURRENT HIV REGIMEN	SWITCHED TO JULUCA
	Weeks 1–48	Weeks 52–100
People who left the study due to side effects	<1%	3%

Of the people who left the study due to side effects, the most common side effects leading to discontinuation were psychiatric disorders:

- 2% of patients who took JULUCA during weeks 1–48
- 2% of patients who took JULUCA during weeks 1–100
- <1% of patients who continued their HIV regimens during weeks 1–48
- 1% of patients who switched to JULUCA during weeks 52–100

*Your results may vary.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA? (cont'd)

JULUCA can cause serious side effects, including: (cont'd)

- **Depression or mood changes.** Tell your healthcare provider right away or get medical help if you have any of the following symptoms:
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself
- **The most common side effects of JULUCA include:**
 - diarrhea
 - headache

These are not all the possible side effects of JULUCA. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider right away if you have any new symptoms while taking JULUCA.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25. Please see [Important Facts](#) about JULUCA on page 27.

Who should not take JULUCA?

Talk with your doctor about all medications you take. **Do not take JULUCA if you take:**

Drug	Examples of Brand Names
Dofetilide	Tikosyn®
Anti-seizure medicines	
carbamazepine	Carbatrol®, Epitol®, Equetro®, Tegretol®, Tegretol®-XR
oxcarbazepine	Oxtellar XR®, Trileptal®
phenobarbital	Luminal®, Solfoton®
phenytoin	Dilantin®, Dilantin-125®, Phenytek®
Anti-tuberculosis medicines	
rifampin	Rifater®, Rifamate®, Rimactane®, Rifadin®
rifapentine	Priftin®
Proton pump inhibitor (PPI) medicines for certain stomach or intestinal problems	
esomeprazole	Nexium®, Vimovo®
lansoprazole	Prevacid®
omeprazole	Prilosec®, Zegerid®
pantoprazole sodium	Protonix®
rabeprazole	AcipHex®, AcipHex® Sprinkle™
St. John's wort (<i>Hypericum perforatum</i>)	
More than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate	

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25. Please see [Important Facts](#) about JULUCA on page 27.

Danny
Undetectable
since 1997



**The years have taught me
the value of asking questions.
Today, I want to ask about JULUCA.**

**IMPORTANT SAFETY INFORMATION
(cont'd)**

Before you take JULUCA, tell your healthcare provider if you:

- have ever had a severe skin rash or an allergic reaction to medicines that contain dolutegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had a mental health problem.
- are pregnant or plan to become pregnant. One of the medicines in JULUCA, called dolutegravir, may harm your unborn baby.
 - Your healthcare provider may prescribe a different medicine than JULUCA if you are planning to become pregnant or if pregnancy is confirmed during the first 12 weeks of pregnancy
 - If you can become pregnant, your healthcare provider will perform a pregnancy test before you start treatment with JULUCA, and you should consistently use effective birth control (contraception) during treatment with JULUCA
 - Tell your healthcare provider right away if you are planning to become pregnant, you become pregnant, or think you may be pregnant during treatment with JULUCA

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25. Please see Important Facts about JULUCA on page 27.

Questions to ask your HIV healthcare team.

To find out if JULUCA may be a next step for you, here's a list of questions:

- 1.** I've heard about JULUCA. What are its benefits and risks?
- 2.** I'm undetectable* now. Is there a reason to consider switching to JULUCA?
- 3.** How many medicines are in my current HIV regimen?
- 4.** How might JULUCA fit in with my health goals?

*Undetectable for at least six months. Undetectable means keeping the amount of HIV-1 in the blood at very low levels (less than 50 copies per mL).

IMPORTANT SAFETY INFORMATION (cont'd)

Before you take JULUCA, tell your healthcare provider if you: (cont'd)

- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take JULUCA.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if JULUCA can pass to your baby in your breast milk.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25. Please see Important Facts about JULUCA on page 27.



Questions to ask yourself.

- 1.** My current HIV treatment is:
- 1 pill each day
 - 2 pills each day
 - more than 2 pills each day

- 2.** My HIV treatment gets in the way of the rest of my life:
- not at all
 - several times a week
 - every day

- 3.** I have side effects:
- never
 - sometimes
 - often

- 4.** I'm undetectable—a once-daily pill with fewer medicines could fit my lifestyle:
- yes
 - no

- 5.** I'm concerned about the effects my HIV medicines may be having on my body:
- not at all
 - sometimes
 - often

IMPORTANT SAFETY INFORMATION (cont'd)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Some medicines interact with JULUCA. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with JULUCA.
- Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take JULUCA with other medicines.

How to take JULUCA?

- **Take JULUCA 1 time a day exactly as your healthcare provider tells you.**
- **Always take JULUCA with a meal.** A protein drink alone does not replace a meal.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.

Access and coverage assistance.



Comprehensive information on access and coverage to help you get your prescribed ViiV Healthcare medicines.

Insured

Dedicated Access Coordinators

ViiVConnect features one-on-one support to assist you in gaining information on coverage and access to ViiV Healthcare medicines.

Underinsured

ViiVConnect Savings Card

The ViiVConnect Savings Card* offers eligible patients savings on additional costs (such as out-of-pocket costs) not covered by insurance.

Uninsured

Patient Assistance Program

The program* offers ViiV Healthcare medicines at no cost to patients who qualify.

See if you're eligible for savings and support:

➤ [Visit viivconnect.com](https://www.viivconnect.com)

➤ [Call 1-844-588-3288 \(toll free\)](tel:1-844-588-3288)

Monday–Friday, 8 AM to 11 PM (ET) | *Multilingual options available*

*Subject to eligibility and program terms and conditions; ViiVConnect programs do not constitute health insurance.

Please see **Important Safety Information** throughout this brochure and a summary on pages 24 and 25.
Please see **Important Facts** about JULUCA on page 27.



 **Juluca**
dolutegravir 50 mg/
rilpivirine 25 mg tablets

**I didn't take any shortcuts to
become the person I am.
And now, there's a path to
stay undetectable with fewer medicines.**

Elveth
Undetectable since 2015

IMPORTANT SAFETY INFORMATION (cont'd)

Do not take JULUCA if you:

- have ever had an allergic reaction to a medicine that contains dolutegravir or rilpivirine.
- are taking any of the following medicines:
 - dofetilide
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifampin
 - rifapentine
 - proton pump inhibitors (including esomeprazole, lansoprazole, omeprazole, pantoprazole sodium, rabeprazole)
 - St. John's wort (*Hypericum perforatum*)
 - more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.

Please see Important Facts about JULUCA on page 27.

Healthy living tips.

➤ **Download the free guide, *Living Healthy With HIV*, now at juluca.com.**

As you stay undetectable, living healthy with HIV becomes a larger focus in your overall health.

Discover how to bring more balance to four areas of your life:

Getting and Giving Support

Discover how to use online communities to strengthen your support system as you take on challenges of living with HIV.

Staying Active

Learn how good it can be to keep your body moving. And learn a few simple exercises to work into your daily routine.

Eating Well

How do vitamins and minerals benefit your body? Find out here—and learn which foods can help support your immune system.

Ways to Unwind

Find out some things you can do to give yourself the attention you deserve.



 **Juluca**
dolutegravir 50 mg/
rilpivirine 25 mg tablets

Reaching undetectable gave me a
huge relief, even with just going
about my day-to-day.

Rodney | Massachusetts

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA?

JULUCA can cause serious side effects, including:

- **Severe skin rash and allergic reactions.** Call your healthcare provider right away if you develop a rash with JULUCA. **Stop taking JULUCA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - blisters or sores in mouth
 - blisters or peeling of the skin
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue
 - problems breathing

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.

Please see [Important Facts](#) about JULUCA on page 27.

Rodney's personal story.

1. Tell us a little bit about your life — how do you like to spend your time?

I try to live somewhat stress-free these days, so I do things that I enjoy. I do some painting and work a bit with my sister to help with her business. I like all kinds of cooking too, and I love trying out different recipes.

2. Reaching undetectable is a major achievement; what was it like for you?

It felt really good, because I had reached a certain point of recovery where I didn't have to get as nervous about getting sick with a cold or something. Overall, I'd say it gave me a huge relief, even with just going about my day-to-day.

3. What's your relationship with your doctor like? When you learn about new HIV treatments, do you discuss them?

In the beginning, I felt like I wasn't getting proper care. Eventually, I found a healthcare group in my city and they were so fantastic. I found another doctor who was also fantastic, and I've been with him ever since. I always ask about new medications when I read something online, and discuss it with my doctor.

4. What have you learned about yourself since you were first diagnosed with HIV?

I learned that I have a lot of resilience. At times I felt like I wouldn't get through it, but I kept on giving myself that little boost...I thought it was fantastic that I had this little warrior inside me.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA? (cont'd)

JULUCA can cause serious side effects, including: (cont'd)

- **Liver problems.** People with a history of hepatitis B or C virus who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during treatment with JULUCA. Liver problems, including liver failure, have also happened in people without history of liver disease or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or "tea-colored" urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain, aching, or tenderness on the right side of your stomach area

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.



 **Juluca**
dolutegravir 50 mg/
rilpivirine 25 mg tablets

I've been through a lot —
but that's the beauty of it.

Jorge | California

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of JULUCA? (cont'd)

JULUCA can cause serious side effects, including: (cont'd)

- **Depression or mood changes. Tell your healthcare provider right away or get medical help if you have any of the following symptoms:**
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself
- **The most common side effects of JULUCA include:**
 - diarrhea
 - headache

These are not all the possible side effects of JULUCA. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider right away if you have any new symptoms while taking JULUCA.

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.

Please see [Important Facts](#) about JULUCA on page 27.

Jorge's personal story.

1. Tell us a little bit about your life — how do you like to spend your time?

I'm a mental health therapist by day and work in an HIV nonprofit organization. I'm also a college professor on the graduate level, and I write for an LGBT magazine. I spend a lot of time with my friends and with my community; I'm a leader and very involved in our LGBT movement.

2. How would friends describe your personality?

Friends I've known for twenty-plus years would say that I'm a fighter and I don't give up. Friends that I've known for five years would say that I do give up. It's not that I do give up, it's just that I know how to pick and choose my battles now. I'm about change and accepting the things that I can't change.

3. What are you most proud of?

I'm proud of a lot of things, but I'm really proud of my parents. I think they're a prime example of change and challenge, culture and religion, being a brown family in America and having to juggle two cultures and so many messages. I am so proud they were able to overcome everything they were taught in order to love and accept their gay, HIV-positive son. And I'm proud of myself, too. I live my life to the fullest without any taboos or stigma. That is what makes me the most proud — being completely free to be who I am.

4. How do you stay healthy — what do you do to stay active?

For me, being healthy is not just physical; it's emotional and mental. I believe in surrounding myself with good, positive people that are going to lift me up. I cannot afford to have negativity, shame, or stigma around me at all. To stay active, I hike — it's really very liberating for me. When I hike, I talk to myself, I cry, I plan, I process, I let go. I'm staying fit, but also healing my soul and my heart.

IMPORTANT SAFETY INFORMATION (cont'd)

Before you take JULUCA, tell your healthcare provider if you:

- have ever had a severe skin rash or an allergic reaction to medicines that contain dolutegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had a mental health problem.
- are pregnant or plan to become pregnant. One of the medicines in JULUCA, called dolutegravir, may harm your unborn baby.
 - Your healthcare provider may prescribe a different medicine than JULUCA if you are planning to become pregnant or if pregnancy is confirmed during the first 12 weeks of pregnancy
 - If you can become pregnant, your healthcare provider will perform a pregnancy test before you start treatment with JULUCA, and you should consistently use effective birth control (contraception) during treatment with JULUCA
 - Tell your healthcare provider right away if you are planning to become pregnant, you become pregnant, or think you may be pregnant during treatment with JULUCA

Please see Important Safety Information throughout this brochure and a summary on pages 24 and 25.
Please see [Important Facts](#) about JULUCA on page 27.

IMPORTANT SAFETY INFORMATION

Do not take JULUCA if you:

- have ever had an allergic reaction to a medicine that contains dolutegravir or rilpivirine.
- are taking any of the following medicines:
 - dofetilide
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifampin
 - rifapentine
 - proton pump inhibitors (including esomeprazole, lansoprazole, omeprazole, pantoprazole sodium, rabeprazole)
 - St. John's wort (*Hypericum perforatum*)
 - more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate

What are the possible side effects of JULUCA?

JULUCA can cause serious side effects, including:

- **Severe skin rash and allergic reactions.** Call your healthcare provider right away if you develop a rash with JULUCA. **Stop taking JULUCA and get medical help right away if you develop a rash with any of the following signs or symptoms:**

- fever
- generally ill feeling
- tiredness
- muscle or joint aches
- blisters or sores in mouth
- blisters or peeling of the skin
- redness or swelling of the eyes
- swelling of the mouth, face, lips, or tongue
- problems breathing

- **Liver problems.** People with a history of hepatitis B or C virus who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during treatment with JULUCA. Liver problems, including liver failure, have also happened in people without history of liver disease or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**

- your skin or the white part of your eyes turns yellow (jaundice)
- dark or "tea-colored" urine
- light-colored stools (bowel movements)
- nausea or vomiting
- loss of appetite
- pain, aching, or tenderness on the right side of your stomach area

- **Depression or mood changes.** Tell your healthcare provider right away or get medical help if you have any of the following symptoms:

- feeling sad or hopeless
- feeling anxious or restless
- have thoughts of hurting yourself (suicide) or have tried to hurt yourself

- **The most common side effects of JULUCA include:**

- diarrhea
- headache

These are not all the possible side effects of JULUCA. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider right away if you have any new symptoms while taking JULUCA.

Please see Important Safety Information throughout this brochure.
Please see [Important Facts](#) about JULUCA on page 27.

IMPORTANT SAFETY INFORMATION (cont'd)

Before you take JULUCA, tell your healthcare provider if you:

- have ever had a severe skin rash or an allergic reaction to medicines that contain dolutegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had a mental health problem.
- are pregnant or plan to become pregnant. One of the medicines in JULUCA, called dolutegravir, may harm your unborn baby.
 - Your healthcare provider may prescribe a different medicine than JULUCA if you are planning to become pregnant or if pregnancy is confirmed during the first 12 weeks of pregnancy
 - If you can become pregnant, your healthcare provider will perform a pregnancy test before you start treatment with JULUCA, and you should consistently use effective birth control (contraception) during treatment with JULUCA
 - Tell your healthcare provider right away if you are planning to become pregnant, you become pregnant, or think you may be pregnant during treatment with JULUCA
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take JULUCA.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if JULUCA can pass to your baby in your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Some medicines interact with JULUCA. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with JULUCA.
- Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take JULUCA with other medicines.

How to take JULUCA?

- **Take JULUCA 1 time a day exactly as your healthcare provider tells you.**
- **Always take JULUCA with a meal.** A protein drink alone does not replace a meal.



Juluca

dolutegravir 50 mg/
rilpivirine 25 mg tablets



For information on access and coverage to help
you get your prescribed ViiV Healthcare medications
Insured • Underinsured • Uninsured

LEARN MORE



1-844-588-3288 (toll free)
Monday-Friday, 8AM-11PM (ET)



www.viivconnect.com

Trademarks are owned by or licensed to the ViiV Healthcare group of companies.

Please see Important Safety Information throughout this
brochure and a summary on pages 24 and 25.
Please see **Important Facts** about JULUCA on page 27.



©2019 ViiV Healthcare or licensor.
DGRWCNT190001 November 2019
Produced in USA.

IMPORTANT FACTS

This is only a brief summary of important information about JULUCA and does not replace talking to your healthcare provider about your condition and treatment.

(Jah-LOO-kah)

ABOUT JULUCA

- JULUCA is a prescription medicine that is used without other antiretroviral medicines to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults to replace their current anti-HIV-1 medicines when their healthcare provider determines that they meet certain requirements.
- HIV-1 is the virus that causes Acquired Immune Deficiency Syndrome (AIDS).
- It is not known if JULUCA is safe and effective in children.

HOW TO TAKE JULUCA

- **Take JULUCA 1 time a day exactly as your healthcare provider tells you.**
- **Always take JULUCA with a meal.** A protein drink alone does not replace a meal.

DO NOT TAKE JULUCA IF YOU

- have ever had an allergic reaction to a medicine that contains dolutegravir or rilpivirine.
- are taking any of the following medicines: dofetilide; carbamazepine; oxcarbazepine; phenobarbital; phenytoin; rifampin; rifapentine; proton pump inhibitors (including esomeprazole, lansoprazole, omeprazole, pantoprazole sodium, rabeprazole); St. John's wort (*Hypericum perforatum*); or more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate.

BEFORE TAKING JULUCA

Tell your healthcare provider if you:

- have ever had a severe skin rash or an allergic reaction to medicines that contain dolutegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had a mental health problem.
- are pregnant or plan to become pregnant. One of the medicines in JULUCA, called dolutegravir, may harm your unborn baby.
 - Your healthcare provider may prescribe a different medicine than JULUCA if you are planning to become pregnant or if pregnancy is confirmed during the first 12 weeks of pregnancy.
 - If you can become pregnant, your healthcare provider will perform a pregnancy test before you start treatment with JULUCA, and you should consistently use effective birth control (contraception) during treatment with JULUCA.
 - Tell your healthcare provider right away if you are planning to become pregnant, you become pregnant, or think you may be pregnant during treatment with JULUCA.
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take JULUCA.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if JULUCA can pass to your baby in your breast milk. Talk with your healthcare provider about the best way to feed your baby.

BEFORE TAKING JULUCA (cont'd)

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that should not be taken with JULUCA.
- Do not start taking a new medicine without telling your healthcare provider.

POSSIBLE SIDE EFFECTS OF JULUCA

JULUCA can cause serious side effects, including:

- **Severe skin rash and allergic reactions.** Call your healthcare provider right away if you develop a rash with JULUCA. **Stop taking JULUCA and get medical help right away if you develop a rash with any of the following signs or symptoms:** fever; generally ill feeling; tiredness; muscle or joint aches; blisters or sores in mouth; blisters or peeling of the skin; redness or swelling of the eyes; swelling of the mouth, face, lips, or tongue; problems breathing.
- **Liver problems.** People with a history of hepatitis B or C virus who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during treatment with JULUCA. Liver problems, including liver failure, have also happened in people without history of liver disease or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:** your skin or the white part of your eyes turns yellow (jaundice); dark or "tea-colored" urine; light-colored stools (bowel movements); nausea or vomiting; loss of appetite; pain, aching, or tenderness on the right side of your stomach area.
- **Depression or mood changes. Tell your healthcare provider right away or get medical help if you have any of the following symptoms:** feeling sad or hopeless; feeling anxious or restless; have thoughts of hurting yourself (suicide) or have tried to hurt yourself.
- **The most common side effects of JULUCA include:** diarrhea and headache.

These are not all the possible side effects of JULUCA. Tell your healthcare provider right away if you have any new symptoms while taking JULUCA. You may report side effects to FDA at 1-800-FDA-1088.

GET MORE INFORMATION

- This is only a brief summary of important information about JULUCA. Talk to your healthcare provider or pharmacist to learn more.
- Go to JULUCA.com or call 1-877-844-8872, where you can also get FDA-approved labeling.



ViiV Healthcare
Research Triangle Park, NC 27709

Trademark is owned by or licensed to the ViiV Healthcare group of companies.